

Research

Platinum-doublet chemotherapy for advanced gastroenteropancreatic neuroendocrine carcinoma: a systematic review and meta-analysis

Akihiro Ohmoto¹  · Yu Fujiwara²  · Nobuyuki Horita³  · Kenji Nakano¹  · Shunji Takahashi¹ 

Received: 14 April 2022 / Accepted: 18 May 2022

Published online: 30 May 2022

© The Author(s) 2022 

Abstract

Background Platinum-doublet chemotherapy has been conventionally used for patients with advanced gastroenteropancreatic (GEP) neuroendocrine carcinoma (NEC) but evidence of chemotherapy is based on studies with small sample sizes and remains scarce. Thus, we conducted a systematic review and meta-analysis to elucidate the efficacy of platinum-doublet chemotherapy for advanced GEP-NEC.

Methods We performed a database search in PubMed/MEDLINE and EMBASE. Eligible studies were prospective and retrospective studies documenting the efficacy of platinum plus etoposide (EP) and platinum plus irinotecan (IP) for advanced GEP-NEC. Overall response rate (ORR), median progression-free survival (PFS), and median overall survival (OS) were pooled and weighted using generic inverse variance in a random-effects meta-analysis model.

Results Nineteen studies including 1157 patients were identified. The ORR of the platinum-doublet regimen, EP, and IP was 49.1% (95% confidence interval [CI], 41.8–56.5), 44.4% (95% CI: 35.9–53.0), and 59.4% (95% CI: 48.0–70.8). The pooled median OS of the platinum-doublet regimen, EP, and IP was 12.9 months (95% CI: 10.9–15.3), 12.9 months (95% CI: 10.8–15.4), and 12.9 months (95% CI: 6.0–27.8), and the pooled median PFS of the platinum-doublet regimen, EP, and IP was 5.4 months (95% CI: 4.5–6.4), 5.4 months (95% CI 4.5–6.5), and 4.0 months (95% CI: 1.4–11.7), respectively.

Conclusion Considerable response rate and survival time of the platinum-doublet regimen for advanced GEP-NEC were observed. IP and EP regimens can be reasonably applicable and these results provide a reference for oncologists in deciding the suitable regimen for patients with advanced GEP-NEC.

Keywords Neuroendocrine carcinoma · Gastroenteropancreatic system · Platinum-doublet · Chemotherapy

Akihiro Ohmoto and Yu Fujiwara contributed equally

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s12672-022-00499-w>.

✉ Akihiro Ohmoto, akihiro.omoto@jfc.or.jp; ✉ Yu Fujiwara, yu.fujiwara@mountsinai.org | ¹Department of Medical Oncology, The Cancer Institute Hospital of Japanese Foundation for Cancer Research, 3-8-31, Ariake, Koto-ku, Tokyo 1358550, Japan. ²Department of Medicine, Icahn School of Medicine at Mount Sinai, Mount Sinai Beth Israel, 281 First Avenue, New York, NY 10003, USA. ³Chemotherapy Center, Yokohama City University Hospital, 3-9 Fukuura, Kanazawa-ku, Yokohama 236-0004, Japan.



1 Introduction

Neuroendocrine carcinoma (NEC) is an aggressive pathological entity of neuroendocrine neoplasms (NENs) and is composed of small cell carcinoma and large cell NEC (LCNEC). According to the 2019 World Health Organization classification of tumors of the digestive system, poorly differentiated tumors with a high mitotic count (> 20 mitoses/10 high-power field) or high Ki-67 index (> 20%) are categorized as NEC [1]. The NEC occurs all over the body, and the most common primary site, except for the lung, is the gastroenteropancreatic (GEP) system. Owing to insufficient preclinical and clinical data, treatment strategies for extrapulmonary NEC are mainly based on those for small cell lung cancer (SCLC), which is based on morphological similarity. The most pervasive initial regimen for advanced cases is platinum-doublet chemotherapy, such as cisplatin (CDDP) plus etoposide (ETP), or CDDP plus irinotecan (CPT-11). To mitigate renal, gastrointestinal, and peripheral nerve toxicities, carboplatin (CBDCA) has become an alternative to CDDP in older and frail patients. The latest National Comprehensive Cancer Network (NCCN) guidelines for neuroendocrine tumors (NETs) are listed in parallel with CDDP/ETP, CBDCA/ETP, CDDP/CPT-11, and CBDCA/CPT-11 as platinum-doublet chemotherapy for unresectable or metastatic extrapulmonary NEC [2]. The selection of these regimens is chiefly based on toxicity profiles, patients' performance status, and their preferences. However, only a few prospective trials have been conducted, and most of the available data were derived from small-scale retrospective studies. Consequently, the reported efficacy of platinum-doublet chemotherapy is heterogeneous among studies, and robust clinical evidence does not exist. To provide helpful information for daily practice, we aimed to conduct a systematic review and meta-analysis of available prospective and retrospective studies evaluating platinum-doublet regimens for GEP-NEC.

2 Materials and methods

2.1 Protocol registration

This systematic review and meta-analysis was performed based on the Preferred Reporting Items for Systematic Reviews and Meta-Analysis criteria. Prospective and retrospective studies that analyzed the efficacy of platinum-containing regimens in patients with advanced GEP-NEC were included in this analysis. The protocol was registered in the PROSPERO registry (CRD42021243614).

2.2 Data sources and searches

Database searches were conducted using the PubMed/MEDLINE and Embase databases. The following keywords were used for literature retrieval: ((neuroendocrine carcinoma) OR (NEC) OR (neuroendocrine tumor) OR (neuroendocrine neoplasm) OR (small cell carcinoma)) AND (((platinum) and (etoposide)) or ((platinum) and (irinotecan)) or ((cisplatin) and (etoposide)) or ((carboplatin) and (etoposide)) or ((cisplatin) and (irinotecan)) or ((carboplatin) and (irinotecan)) or (platinum-based)). We searched these databases on July 07, 2021. Two authors (AO and YF) independently performed additional manual searches.

2.3 Selection criteria

Eligible studies included randomized trials, single-arm trials, prospective observational studies, and retrospective studies on the efficacy of platinum-based chemotherapy for advanced NEC. Case reports and case series with a sample size of three or fewer cases were excluded because of insufficient evaluation of the efficacy of chemotherapy. All the eligible articles were written in English.

We selected advanced or inoperable patients pathologically diagnosed with NEC using the GEP system to conduct a comprehensive review of platinum-based chemotherapy for these populations. Reports of well-differentiated NETs were excluded.

Studies evaluating either a platinum plus ETP regimen (EP) or a platinum plus CPT-11 regimen (IP) were included in the analysis. Platinum-based regimens with different doses and densities were regarded the same because of the rarity of NEC. Reports evaluating chemotherapy in the neoadjuvant or adjuvant settings were excluded.

2.4 Data extraction and risk of bias assessment

Two authors independently extracted data from eligible studies (AO and YF). Study characteristics included the name of the first author, title of each study, publication year, number of participants, sex, age, clinical stage, primary site, chemotherapy regimen, dosage and intensity of chemotherapy, hazard ratio (HR) of overall survival (OS) and progression-free survival (PFS), survival ratio of OS and PFS at a specific point, median OS and PFS with 95% confidence interval (95% CI), overall response rate (ORR), and complete response rate (CRR). Risk of bias was assessed using the 9-star Ottawa Newcastle Scale for non-randomized studies and the Cochrane risk of bias tool for randomized controlled trials.

2.5 Data synthesis and statistical analysis

The pooled ORR, CRR, median PFS, and median OS were calculated using a random-effects meta-analysis model with the generic inverse variance method. Agresti-Coull adjustment was applied to estimate the standard error of the binomial proportion [3]. RevMan 5.4 was used for calculating these data [4]. The statistical significance threshold was set at $p < 0.05$. A two-tailed level of 0.10 and $I^2 < 50\%$ were set for heterogeneity.

3 Results

3.1 Study selection

By searching the databases, 10,591 articles were found, 3279 articles were removed due to duplicates, and 7310 studies underwent further screening. Reports on lung cancer ($n = 4208$) were removed during this screening process, and 165 articles were potentially eligible after screening for the abstracts and titles. Of these studies, 146 reports were excluded, and 19 reports that included 1,157 patients were finally identified [5–23]. A flowchart of the study selection process is shown in Fig. 1.

3.2 Study characteristic and treatment group description

Among the 19 studies, three directly compared the IP and EP regimens, of which one was a randomized phase II trial and the other two were retrospective studies [5, 6, 23]. The other 16 studies included one prospective study evaluating EP [15] and 15 retrospective single-arm studies, among which four studies analyzed IP and 11 studies evaluated EP. All the studies evaluated patients with advanced GEP-NEC who received systemic chemotherapy (IP or EP). Key study characteristics are summarized in Table 1. Among 430 patients whose primary organs were reported, the most common was the esophagus (29%), stomach (25%), and pancreas (22%). Information regarding the chemotherapy setting was available in 17 studies (Table 1). Among these studies, 14 included only chemo-naïve patients in the first-line setting. Regarding the remaining three studies, Zhang et al. evaluated the first-line treatment or adjuvant chemotherapy, Lu et al. included 15 patients in the first-line setting and one patient in the later-line, and Pulvirenti reported that 97% of included patients were treated with platinum-chemotherapy in the first-line setting [5, 9, 23]. 9-star Ottawa Newcastle Scale was used to evaluate the risk of bias of the included 18 studies, and the Cochrane risk bias assessment was used for one randomized phase II trial (Supplementary Table 1).




Fig. 1 Flowchart of the systematic review process. On the basis of the initial screening and eligibility assessment, 19 studies with 1157 patients were identified

3.3 Meta-analysis of response rate

We evaluated ORR and CRR in patients with NEC who received EP and IP. For ORR, 15 studies that evaluated 987 patients treated with EP and seven studies that included 377 patients treated with IP were included. The pooled ORR in all patients who received the platinum-doublet regimen was 49.1% (95% CI 41.8–56.5%), and subgroup analysis for the EP and IP groups showed ORR of 44.4% (95% CI 35.9–53.0%) and 59.4% (95% CI 48.0–70.8%), respectively (Fig. 2). The CDDP/ETP (seven studies) and CBDCA/ETP (three studies) groups showed ORR of 31.0% (95% CI 24.3–37.6%) and 67.2% (95% CI 18.0–116.3%), respectively. For 370 patients in six studies who received CDDP/CPT-11, the pooled ORR was 58.1% (95% CI 46.1–70.0%) (Supplementary Fig. 1). Three studies compared the efficacy of EP and IP directly, and the odds ratio of ORR using EP as a reference was 1.95 (95% CI 0.86–4.39%, $p = 0.11$) (Fig. 3).

Regarding CRR, 10 studies with 391 participants evaluating EP and four studies with 45 patients assessing IP were included. The pooled CRR in all patients who received the platinum-doublet regimen was 2.1% (95% CI – 0.3% to 4.5%), and subgroup analysis for the EP and IP groups showed CRR of 2.1% (95% CI – 1.0% to 5.2%) and 5.8% (95% CI – 3.9% to 15.5%), respectively (Fig. 4). The CDDP/ETP and CBDCA/ETP groups showed CRRs of 1.6% (95% CI – 0.8%

Table 1 Characteristics of selected studies

Authors [Reference]	Study design	Regimen	Treatment setting	Num-ber of patients	Age (median)	Primary organ	Detailed primary organ
Zhang [5]	Randomized phase II	CDDP + ETP CDDP + CPT-11	Chemo-naïve or treated as adjuvant chemotherapy	33 33	NA NA	GEP (N = 58), unknown primary (N = 8)	Stomach (N = 20), Esophagus (N = 13), Colorectum (N = 11), Unknown primary (N = 8), Pancreas (N = 7), Duodenum (N = 4), Small intestine (N = 3)
Yamaguchi [6]	Retrospective	CDDP + ETP CDDP + CPT-11	Chemo-naïve	46 160	NA NA	GEP	Esophagus (N = 75), Stomach (N = 58), Pancreas (N = 29), Hepato-biliary tract (N = 23), Colorectum (N = 17), Small bowel (N = 4)
Sorbye [7]	Retrospective	CDDP + ETP CBDCA + ETP	Chemo-naïve	129 67	NA NA	GEP, unknown primary	NA
Okuma [8]	Retrospective	CDDP + CPT-11	Chemo-naïve	12	62	GEP	Esophagus (N = 12)
Lu [9]	Retrospective	CDDP + CPT-11	Chemo-naïve (N = 15) or treated (N = 1)	16	57	GEP (N = 15), unknown primary (N = 1)	Stomach (N = 8), Esophagus (N = 5), Small intestine (N = 1), Pancreas (N = 1), Unknown primary (N = 1)
Iwasa [10]	Retrospective	CDDP + ETP	Chemo-naïve	21	57	GEP	Pancreas (N = 10), Gallbladder (N = 8), Liver (N = 2), Ampulla of Vater (N = 1)
Okita [11]	Retrospective	CDDP + CPT-11	NA *	12	62	GEP	Stomach (N = 12)
Chin [12]	Retrospective	CDDP + CPT-11	Chemo-naïve	12	66	GEP	Esophagus (N = 12)
Yoon [13]	Retrospective	CDDP + ETP	Chemo-naïve	64	57	GEP	Gastrointestinal tract (N = 31), Hepatobiliary tract (N = 25), Abdominal lymph node (N = 8)
Patta [14]	Retrospective	CDDP + ETP	Chemo-naïve	8	64	GEP	Colorectum (N = 8)
Walter [15]	Prospective cohort	Platinum + ETP	Chemo-naïve	152	NA	GEP, unknown primary	NA
Brandi [16]	Retrospective	Platinum + ETP	Chemo-naïve	21	58	GEP (N = 16), unknown primary (N = 5)	Pancreas (N = 10), unknown primary (N = 5), stomach (N = 2), colon (N = 2), duodenum (N = 1), gallbladder (N = 1)
Heetfeld [17]	Retrospective	Platinum + ETP	Chemo-naïve	113	NA	GEP	NA
Bongiovanni [18]	Retrospective	Platinum + ETP	Chemo-naïve	20	60	GEP	Stomach (N = 8), pancreas (N = 7), colorectum (N = 5)
Hudson [19]	Retrospective	CBDCA + ETP	NA *	6	69	GEP	Esophagus (N = 6)
Sakamoto [20]	Retrospective	CBDCA + ETP	Chemo-naïve	4	66.5	GEP	Pancreas (N = 4)
Kim [21]	Retrospective	CDDP + ETP	Chemo-naïve	17	NA	GEP	NA
Gerard [22]	Retrospective	Platinum + ETP	Chemo-naïve	24	NA	GEP, unknown primary	NA

Table 1 (continued)

Authors [Reference]	Study design	Regimen	Treatment setting	Num-ber of patients	Age (median)	Primary organ	Detailed primary organ
Pulvirenti [23]	Retrospective	Platinum + ETP Platinum + CPT-11	Chemo-naïve **	22 4	NA NA	GEP	Pancreas (N = 26)

CDDP cisplatin, *CPT-11* irinotecan, *ETP* etoposide, *GEP* gastroenteropancreatic, *NA* not available

*Although detailed treatment settings were not clearly documented in the article, the information from the literature suggests the 1st-line setting treatment

** 97% of patients received platinum-based chemotherapy in the 1st-line setting




Fig. 2 Pooled ORR in patients treated with platinum-doublet chemotherapy. 15 studies that evaluated patients treated with EP and seven studies that evaluated patients with IP were included in this analysis. ORR overall response rate, EP platinum plus etoposide, IP platinum plus irinotecan




Fig. 3 Odds ratio of ORR using EP as reference. Three studies that directly compared the ORRs of IP and EP were included in this analysis. Odds ratios were calculated using EP as a reference. Among them, one was a randomized phase II trial (5) and the other two were retrospective studies (6, 23). EP platinum plus etoposide, IP platinum plus irinotecan, ORR overall response rate

to 4.0%) and 12.0% (95% CI – 11.0% to 35.1%), respectively (Supplementary Fig. 2). Studies directly comparing the CRR of the EP regimen with that of the IP regimen were not available.

3.4 Meta-analysis of survival time

Only one study that directly compared EP and IP documented the hazard ratio (HR) of OS. OS and PFS data at a specific time were insufficient in the included studies; therefore, median OS and PFS were integrated to assess the efficacy of the EP and IP regimens (Table 1). The pooled median OS in all patients who received the platinum-doublet regimen




Fig. 4 Pooled CRR in patients treated with platinum-doublet chemotherapy. 10 studies that evaluated patients treated with EP and four studies that evaluated patients treated with IP were included in this analysis. CRR complete response rate, EP platinum plus etoposide, IP platinum plus irinotecan




Fig. 5 Pooled median OS in patients treated with platinum-doublet chemotherapy. Five studies that evaluated patients treated with EP and two studies that evaluated patients with IP were included in this analysis. EP platinum plus etoposide, IP platinum plus irinotecan, OS overall survival




Fig. 6 Pooled median PFS in patients treated with platinum-doublet chemotherapy. Eight studies that evaluated patients treated with EP and one study that evaluated patients with IP were included in this analysis. *EP* platinum plus etoposide, *IP* platinum plus irinotecan, *PFS* progression-free survival

was 12.9 months (95% CI 10.9–15.3 months), whereas the EP and IP subgroups had the median OS of 12.9 months (95% CI 10.8–15.4 months) and 12.9 months (95% CI 6.0–27.8 months), respectively (Fig. 5). The Median OS of the subgroups based on each regimen is shown in Supplementary Fig. 3. The pooled median PFS was 5.4 months (95% CI 4.5–6.4 months) in all patients with platinum-regimen, 5.4 months (95% CI 4.5–6.5 months) in the EP subgroup, and 4.0 months (95% CI 1.4–11.7 months) in the IP subgroup (Fig. 6). The Median PFS of the subgroups based on each regimen is shown in Supplementary Fig. 4.

4 Discussion

To the best of our knowledge, this is the first systematic review and meta-analysis of the clinical efficacy of platinum-doublet chemotherapy for advanced GEP-NEC. Most of the clinical data about this rare malignancy are derived from retrospective analyses, and only one randomized phase II trial data have been published to date [5]. Morizane et al. recently presented phase III trial (JCOG 1213) data between CDDP/ETP and CDDP/CPT-11 in 170 patients with GEP-NEC [24]. Our large-scale analysis, including more than 1100 patients, presents referential information about suitable chemotherapy options for GEP-NEC. According to the Surveillance, Epidemiology, and End Results database analysis including more than 160,000 NEC cases, the most common primary organ, except for the lung, was the GEP system, and the ratio of GEP-primary NEC to all extrapulmonary NEC cases was 37% [25]. In our analysis, studies that included extrapulmonary NEC other than GEP-NEC (e.g., urological or gynecological NEC) were excluded. Current treatment strategies generally differ between tumors in the GEP system and those in genitourinary or gynecological systems, and no uniform guidelines exist for NEC all over the body.

In our analysis, the pooled ORR in all patients with the platinum-doublet regimen was 49%, and subgroup analysis for the EP and IP groups exhibited ORRs of 44% and 59%, respectively. Three comparative studies between EP and IP did not detect significant differences in the ORR. These pooled median OS and PFS were similar between the two groups (OS: 12.9 months for EP and 12.9 months for IP; PFS: 5.4 months for EP and 4.0 months for IP). Although our analysis showed a favorable pooled ORR in EP and IP, the median PFS and OS of both treatments were short.

In the JCOG 1213 trial, no significant difference was observed in OS or PFS between the EP and IP groups (median OS 12.5 months vs. 10.9 months; HR of EP to IP 1.04, 90% CI 0.79–1.37; median PFS 5.6 months vs. 5.1 months; HR 1.06, 95% CI 0.78–1.45). The ORRs were 54.5% and 52.5% in the EP and IP groups, respectively. These outcomes were generally consistent with those of our analysis. Several trials have demonstrated OS prolongation by adding anti-programmed death ligand 1 antibodies (atezolizumab or durvalumab) to EP, and a similar approach may be promising for GEP-NEC [26, 27]. The European Neuroendocrine Tumor Society guidelines state that CDDP/ETP is a standard regimen for GEP-NEC, and CPT-11 could be an alternative to ETP [28]. According to the European Society of Medical Oncology clinical practice guidelines for the treatment of advanced GEP-NENs, CDDP or CBDCA plus ETP is regarded as a standard first-line regimen for NEC [29]. Although the number of comparative studies between the regimens is limited, our results suggest no definite priority between EP and IP. Another interesting finding is the seemingly high ORR of CBDCA/ETP (67.2%) compared to that of CDDP/ETP (31.0%). It should be noted that the sample size in the CBDCA/ETP group was less than 100 in only three studies, and no comparative data were available between the two regimens. Therefore, we cannot make a definite comment on the priority of CBDCA/ETP over CDDP/ETP. However, CBDCA might be a better choice when patients are ineligible for CDDP due to the poor general condition or organ dysfunction. Our data support that CBDCA can be a reasonable substitute for CDDP.

For advanced SCLC, a direct comparison of CDDP plus ETP versus CDDP plus CPT-11 is available through several randomized clinical trials and meta-analyses. A phase III trial in Japan comparing CDDP/ETP and CDDP/CPT-11 demonstrated higher ORR and longer OS in patients treated with CDDP/CPT-11 (ORR 67.5%, 95% CI 55.9–77.8% vs. 84.4%, 95% CI 74.4–91.7%, $P=0.02$; median OS 9.4 months vs. 12.8 months, $P=0.002$; HR 0.60, 95% CI: 0.43–0.83) [30]. However, subsequent trials in the United States could not validate the superiority of CDDP/CPT-11 over CDDP/ETP for the ORR or OS [31, 32]. Two independent meta-analyses of randomized trials comparing the efficacy of EP and IP in patients with advanced SCLC showed an OS benefit in the IP group (Jiang et al., pooled HR 0.81, 95% CI: 0.66–0.99, $P=0.044$; Lima et al. pooled HR 0.87, 95% CI: 0.78–0.97, $P=0.02$) [33, 34]. The latest NCCN guidelines for SCLC refer to CDDP/ETP with durvalumab and CBDCA/ETP with atezolizumab or durvalumab as the preferred regimens [35]. The situation is more controversial in extrapulmonary NEC, including GEP-NEC.

Essentially, it is debatable whether chemotherapy response or prognosis is different between SCLC and extrapulmonary NEC. Terashima et al. retrospectively compared 95 SCLC cases and 41 extrapulmonary NEC cases, including 18 with gastrointestinal and 16 with hepatobiliary/pancreatic (HBP) NEC treated with a platinum-based regimen and reported that extrapulmonary NEC had a lower ORR than SCLC (78% vs. 31%, $P<0.01$) [36]. Dasari et al. compared the prognosis of patients with extrapulmonary NEC to those with SCLC and reported that colon, pancreas, and liver NEC had worse OS (HR 1.09, 95% CI: 1.03–1.16; 1.10, 95% CI: 1.03–1.18; 1.85, 95% CI: 1.57–2.18) compared with SCLC, whereas small intestine NEC had better OS (HR 0.56, 95% CI: 0.49–0.64) than SCLC [25]. As a premise, the ORR in patients with SCLC from the aforementioned phase III clinical trials ranges from 44 to 68% in EP and from 48 to 84% in IP, and clinicians should consider such heterogeneous responses among studies comparing the efficacy of chemotherapy for SCLC and GEP-NEC [30–32].

Our study has several limitations. First, as one of the large faults, all other studies except for two studies included in this analysis were retrospective in nature. Second, safety analysis was not performed because of a lack of information in the studies included in this meta-analysis. Third, subgroup analysis between HBP-NEC and GI-NEC or SCLC and LCNEC could not be performed. As Yamaguchi et al. pointed out the distribution between HBP-NEC and GI-NEC affects the clinical efficacy of chemotherapy because clinical behaviors are originally worse in patients with HBP-NEC than in those with GI-NEC [6]. This limitation needs to be addressed by conducting a larger multi-institutional and international clinical trial or by performing an individual patient data meta-analysis. Fourth, the results of CRR obtained from our analysis were highly heterogeneous partly due to a limited number of available studies. Therefore, caution should be applied when interpreting results. Finally, to simply assess the efficacy of chemotherapy in patients with GEP-NEC, our analysis excluded several studies that categorized patients with both GEP-NEC and non-GEP NEC as “patients with NEC.”

In conclusion, our novel systematic review and meta-analysis to assess the efficacy of platinum-doublet chemotherapy in patients with GEP-NEC showed considerable ORR. Subgroup analysis, including three comparative studies between EP and IP, did not detect any significant ORR differences. In terms of efficacy, there is no definite superiority between EP and IP, and both regimens are reasonably applicable to patients with GEP-NEC. We believe that these results provide reference information to clinicians for choosing suitable chemotherapy for this rare malignancy. Larger randomized clinical trials comparing these regimens and individual patient data meta-analyses integrating each patient’s clinical information are

needed to stratify “good responders” to each treatment and identify predictive factors for each therapy, which provides robust clinical evidence for patients with GEP-NEC.

Acknowledgements We would like to thank Editage (www.editage.com) for English language editing.

Author contributions AO, YF: conception and design, collection and synthesis, drafting of the article. NH: statistical analysis of data and interpretation of results. KN, ST: critical revision of the article for important intellectual content and final approval of the article. All authors read and approved the final manuscript.

Funding The authors have no funding for any research relevant to this study.

Data Availability All data generated or analyzed during this study are included in this article. Further inquiries can be directed to the corresponding author.

Declarations

Ethics approval and consent to participate An ethics statement is not applicable because this study is based exclusively on published literature.

Competing interests The authors have no conflicts of interest to declare.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Nagtegaal ID, Odze RD, Klimstra D, et al. The 2019 WHO classification of tumours of the digestive system. *Histopathology*. 2020;76:182–8.
2. National Comprehensive Cancer Network. Neuroendocrine and Adrenal Tumors (Version 4.2021). [cited December 14, 2021]. https://www.nccn.org/professionals/physician_gls/pdf/neuroendocrine.pdf
3. Agresti A, Coull BA. Approximate is better than “exact” for interval estimation of binomial proportions. *Am Stat*. 1998;52:119–26.
4. The Cochrane Collaboration. Review manager (RevMan) [Computer program]. Version 5.4. 2020.
5. Zhang P, Li J, Li J, et al. Etoposide and cisplatin versus irinotecan and cisplatin as the first-line therapy for patients with advanced, poorly differentiated gastroenteropancreatic neuroendocrine carcinoma: a randomized phase 2 study. *Cancer*. 2020;126(Suppl 9):2086–92.
6. Yamaguchi T, Machida N, Morizane C, et al. Multicenter retrospective analysis of systemic chemotherapy for advanced neuroendocrine carcinoma of the digestive system. *Cancer Sci*. 2014;105:1176–81.
7. Sorbye H, Welin S, Langer SW, et al. Predictive and prognostic factors for treatment and survival in 305 patients with advanced gastrointestinal neuroendocrine carcinoma (WHO G3): the NORDIC NEC study. *Ann Oncol*. 2013;24:152–60.
8. Okuma HS, Iwasa S, Shoji H, et al. Irinotecan plus cisplatin in patients with extensive-disease poorly differentiated neuroendocrine carcinoma of the esophagus. *Anticancer Res*. 2014;34:5037–41.
9. Lu ZH, Li J, Lu M, et al. Feasibility and efficacy of combined cisplatin plus irinotecan chemotherapy for gastroenteropancreatic neuroendocrine carcinomas. *Med Oncol*. 2013;30:664.
10. Iwasa S, Morizane C, Okusaka T, et al. Cisplatin and etoposide as first-line chemotherapy for poorly differentiated neuroendocrine carcinoma of the hepatobiliary tract and pancreas. *Jpn J Clin Oncol*. 2010;40:313–8.
11. Okita NT, Kato K, Takahari D, et al. Neuroendocrine tumors of the stomach: chemotherapy with cisplatin plus irinotecan is effective for gastric poorly-differentiated neuroendocrine carcinoma. *Gastric Cancer*. 2011;14:161–5.
12. Chin K, Baba S, Hosaka H, et al. Irinotecan plus cisplatin for therapy of small-cell carcinoma of the esophagus: report of 12 cases from single institution experience. *Jpn J Clin Oncol*. 2008;38:426–31.
13. Yoon SE, Kim JH, Lee SJ, et al. The impact of primary tumor site on outcomes of treatment with etoposide and cisplatin in grade 3 gastroenteropancreatic neuroendocrine carcinoma. *J Cancer*. 2019;10:3140–4.
14. Patta A, Fakhri M. First-line cisplatin plus etoposide in high-grade metastatic neuroendocrine tumors of colon and rectum (MCRC NET): review of 8 cases. *Anticancer Res*. 2011;31:975–8.
15. Walter T, Tougeron D, Baudin E, et al. Poorly differentiated gastro-entéro-pancreatic neuroendocrine carcinomas: are they really heterogeneous? Insights from the FFCD-GTE national cohort. *Eur J Cancer*. 2017;79:158–65.
16. Brandi G, Paragona M, Campana D, et al. Good performance of platinum-based chemotherapy for high-grade gastroenteropancreatic and unknown primary neuroendocrine neoplasms. *J Chemother*. 2018;30:53–8.
17. Heetfeld M, Chougnat CN, Olsen IH, et al. Characteristics and treatment of patients with G3 gastroenteropancreatic neuroendocrine neoplasms. *Endocr Relat Cancer*. 2015;22:657–64.
18. Bongiovanni A, Riva N, Ricci M, et al. First-line chemotherapy in patients with metastatic gastroenteropancreatic neuroendocrine carcinoma. *Onco Targets Ther*. 2015;8:3613–9.

19. Hudson E, Powell J, Mukherjee S, et al. Small cell oesophageal carcinoma: an institutional experience and review of the literature. *Br J Cancer*. 2007;96:708–11.
20. Sakamoto H, Kitano M, Komaki T, et al. Small cell carcinoma of the pancreas: role of EUS-FNA and subsequent effective chemotherapy using carboplatin and etoposide. *J Gastroenterol*. 2009;44:432–8.
21. Kim HK, Ha SY, Lee J, et al. The impact of pathologic differentiation (well/poorly) and the degree of Ki-67 index in patients with metastatic WHO grade 3 GEP-NECs. *Oncotarget*. 2017;8:73974–80.
22. Gerard L, Garcia J, Gauthier A, et al. ctDNA in neuroendocrine carcinoma of gastroenteropancreatic origin or of unknown primary: the CIRCAN-NEC pilot study. *Neuroendocrinology*. 2021;111:951–64.
23. Pulvirenti A, Raj N, Cingarlini S, et al. Platinum-based treatment for well- and poorly differentiated pancreatic neuroendocrine neoplasms. *Pancreas*. 2021;50:138–46.
24. Morizane C, Machida N, Honma Y, et al. Randomized phase III study of etoposide plus cisplatin versus irinotecan plus cisplatin in advanced neuroendocrine carcinoma of the digestive system: a Japan Clinical Oncology Group study (JCOG1213). *J Clin Oncol*. 2022;40:501–501.
25. Dasari A, Mehta K, Byers LA, et al. Comparative study of lung and extrapulmonary poorly differentiated neuroendocrine carcinomas: a SEER database analysis of 162,983 cases. *Cancer*. 2018;124:807–15.
26. Horn L, Mansfield AS, Szczesna A, et al. First-line atezolizumab plus chemotherapy in extensive-stage small-cell lung cancer. *N Engl J Med*. 2018;379:2220–9.
27. Paz-Ares L, Dvorkin M, Chen Y, et al. Durvalumab plus platinum-etoposide versus platinum-etoposide in first-line treatment of extensive-stage small-cell lung cancer (CASPIAN): a randomised, controlled, open-label, phase 3 trial. *Lancet*. 2019;394:1929–39.
28. Garcia-Carbonero R, Sorbye H, Baudin E, et al. ENETS consensus guidelines for high-grade gastroenteropancreatic neuroendocrine tumors and neuroendocrine carcinomas. *Neuroendocrinology*. 2016;103:186–94.
29. Pavel M, Öberg K, Falconi M, et al. Gastroenteropancreatic neuroendocrine neoplasms: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2020;31:844–60.
30. Noda K, Nishiwaki Y, Kawahara M, et al. Irinotecan plus cisplatin compared with etoposide plus cisplatin for extensive small-cell lung cancer. *N Engl J Med*. 2002;346:85–91.
31. Hanna N, Bunn PA Jr, Langer C, et al. Randomized phase III trial comparing irinotecan/cisplatin with etoposide/cisplatin in patients with previously untreated extensive-stage disease small-cell lung cancer. *J Clin Oncol*. 2006;24:2038–43.
32. Lara PN Jr, Natale R, Crowley J, et al. Phase III trial of irinotecan/cisplatin compared with etoposide/cisplatin in extensive-stage small-cell lung cancer: clinical and pharmacogenomic results from SWOG S0124. *J Clin Oncol*. 2009;27:2530–5.
33. Jiang J, Liang X, Zhou X, et al. A meta-analysis of randomized controlled trials comparing irinotecan/platinum with etoposide/platinum in patients with previously untreated extensive-stage small cell lung cancer. *J Thorac Oncol*. 2010;5:867–73.
34. Lima JP, dos Santos LV, Sasse EC, et al. Camptothecins compared with etoposide in combination with platinum analog in extensive stage small cell lung cancer: systematic review with meta-analysis. *J Thorac Oncol*. 2010;5:1986–93.
35. National Comprehensive Cancer Network. Small cell lung cancer (version 2.2022). [cited November 24, 2021]. https://www.nccn.org/professionals/physician_gls/pdf/sclc.pdf
36. Terashima T, Morizane C, Hiraoka N, et al. Comparison of chemotherapeutic treatment outcomes of advanced extrapulmonary neuroendocrine carcinomas and advanced small-cell lung carcinoma. *Neuroendocrinology*. 2012;96:324–32.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.